

Assessment of Knowledge, Attitudes, and Practices Related to Diabetes Mellitus among Patients in Primary Healthcare Settings: A Cross-Sectional Study

Muhammad Tahir¹, Hajra Malik²

1. The University of Lahore, Lahore, Pakistan

2. Riphah International University, Pakistan

* Correspondence: hajimalik87@gmail.com

Received: 08 January 2026 | Revised: 13 February 2026 | Accepted: 28 May 2026 | Published: 30 June 2026

Copyright: © 2026 The Authors. | Publisher: Therapy Plus Clinics, Pakistan

License: This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) License.

ABSTRACT

Background: Diabetes mellitus is a major public health concern that requires active patient participation in treatment and self-management. Adequate knowledge and positive attitudes are essential for promoting appropriate self-care practices and preventing diabetes-related complications. However, evidence regarding diabetes-related knowledge, attitudes, and practices among patients receiving care at primary healthcare facilities in Pakistan remains limited. **Objective:** To assess knowledge, attitudes, and practices regarding diabetes mellitus and identify predictors of good self-care practices among patients attending primary healthcare settings in Lahore, Pakistan. **Methods:** A cross-sectional observational study was conducted among 180 adult patients with diabetes attending primary healthcare facilities in Lahore. Data were collected using a structured, interviewer-administered questionnaire covering sociodemographic characteristics, diabetes-related knowledge, attitudes toward disease management, and self-care practices. Descriptive statistics, chi-square tests, and multivariable logistic regression were used for data analysis. A p-value of <0.05 was considered statistically significant. **Results:** Of the 180 participants, 54.4% had adequate knowledge of diabetes, 62.8% demonstrated positive attitudes toward diabetes management, and 41.7% reported good self-care practices. Higher educational attainment was significantly associated with adequate knowledge (OR 3.12; 95% CI 1.46–6.66; $p=0.003$). Participants with adequate knowledge (OR 3.47; 95% CI 1.84–6.55; $p<0.001$) and positive attitudes (OR 4.18; 95% CI 2.17–8.05; $p<0.001$) were more likely to demonstrate good practices. Medication adherence was reported by 68.9% of participants, whereas adherence to a recommended diabetic diet and regular physical activity was reported by only 39.4% and 36.1%, respectively. In the adjusted analysis, adequate knowledge (adjusted OR 2.96; 95% CI 1.49–5.89; $p=0.002$), positive attitudes (adjusted OR 3.41; 95% CI 1.72–6.77; $p<0.001$), and higher education (adjusted OR 2.18; 95% CI 1.07–4.45; $p=0.031$) independently predicted good self-care practices. **Conclusion:** Although more than half of the participants had adequate diabetes-related knowledge and positive attitudes, self-care practices remained suboptimal, particularly regarding diet and physical activity. Structured diabetes education, behavioural counseling, and patient-centered support should be integrated into primary healthcare services to improve diabetes self-management.

Keywords: Attitudes; Diabetes mellitus; Knowledge; Primary healthcare; Self-care practices; Pakistan

INTRODUCTION

Diabetes mellitus is a chronic metabolic disorder characterized by persistent hyperglycemia resulting from defects in insulin secretion, insulin action, or both, and it represents one of the leading causes of cardiovascular disease, renal failure, blindness, and premature mortality worldwide (1). The global burden of diabetes continues to rise rapidly, particularly in low- and middle-income countries, where health systems face increasing pressure from long-term disease management and diabetes-related complications (2). Effective control of diabetes depends not only on pharmacological treatment but also on patients' active participation in self-care behaviors, making patient-centered management a cornerstone of diabetes control strategies (3).

Knowledge, attitude, and practices (KAP) related to diabetes mellitus are widely recognized as critical determinants of disease outcomes. Adequate knowledge enables patients to understand disease mechanisms, recognize symptoms, adhere to medications, and adopt preventive behaviors, while positive attitudes influence

motivation, treatment compliance, and engagement with health care services (4). Conversely, inappropriate practices such as poor dietary habits, physical inactivity, and irregular monitoring of blood glucose levels contribute to poor glycemic control and accelerate the development of complications (5). Evidence from previous studies suggests that even when patients possess reasonable knowledge about diabetes, this does not consistently translate into optimal self-care practices, indicating a complex interplay between cognitive, behavioral, and contextual factors (6).

In South Asia, including Pakistan, the prevalence of diabetes mellitus has increased substantially due to rapid urbanization, sedentary lifestyles, unhealthy dietary patterns, and population aging (7). Pakistan is currently among the countries with the highest number of adults living with diabetes, and projections indicate a continued rise in prevalence over the coming decades (8). Despite this growing burden, diabetes-related complications remain common, largely due to delayed diagnosis, limited access to structured education, and suboptimal self-management practices among patients (9). Primary health care settings serve as the first and most frequent point of contact for patients with diabetes and therefore play a pivotal role in patient education, counseling, and long-term disease monitoring.

Previous KAP studies on diabetes have demonstrated considerable variation in patient knowledge levels, attitudes, and self-care behaviors across different populations and health care settings (10,11). However, much of the available evidence from Pakistan has focused on tertiary care hospitals or community-based surveys, with comparatively limited data derived from primary health care facilities, where the majority of patients receive routine diabetes care (12). Moreover, there is a lack of integrated assessment examining knowledge, attitude, and practices simultaneously among patients attending primary health care settings in large urban centers such as Lahore. This gap limits the ability of health planners and clinicians to design targeted educational interventions tailored to primary care populations.

Understanding the existing levels of diabetes-related knowledge, attitudes, and practices among patients attending primary health care facilities is essential for identifying gaps in patient education and for strengthening self-management support within routine care. Therefore, the objective of this study was to assess the knowledge, attitude, and practices regarding diabetes mellitus among patients attending primary health care settings in Lahore using a cross-sectional study design (13).

MATERIALS AND METHODS

A cross-sectional observational study was conducted in primary health care settings in Lahore, Pakistan, over a defined data collection period. This study design was selected to assess the prevailing levels of knowledge, attitude, and practices (KAP) regarding diabetes mellitus among patients at a single point in time, which is appropriate for identifying gaps in awareness and self-care behaviors within routine primary care populations (14). Adult patients were approached consecutively during regular clinic hours to minimize selection related to appointment timing. All eligible participants were informed about the study objectives and procedures, and written informed consent was obtained prior to participation.

The study population comprised adult patients aged 18 years and above who were attending primary health care facilities for diabetes-related or general medical consultations. Patients with a confirmed diagnosis of diabetes mellitus or those receiving diabetes-related counseling were included. Individuals who were severely ill, cognitively impaired, or unwilling to participate were excluded to ensure reliable responses. A non-probability convenience sampling technique was employed due to feasibility constraints inherent to busy primary care settings; however, recruitment across multiple centers and different days was undertaken to enhance sample heterogeneity and reduce selection bias. Data were collected using a structured, interviewer-administered questionnaire developed based on previously published and widely used diabetes KAP instruments (15,16). The questionnaire consisted of four sections: socio-demographic characteristics, knowledge related to diabetes mellitus, attitude toward diabetes management, and self-care practices. The knowledge domain assessed understanding of disease causes, symptoms, complications, and preventive measures. The attitude domain evaluated beliefs, perceptions, and motivation regarding diabetes control and treatment adherence. The practices domain assessed routine behaviors including dietary adherence, physical activity, medication compliance, blood glucose monitoring, and follow-up visits. Responses were scored using predefined criteria, and overall KAP levels were categorized as adequate or inadequate knowledge, positive or negative attitude, and good or poor practices based on established cut-off values used in similar studies (17).

Several measures were implemented to reduce information bias and enhance data quality. Interviews were conducted by trained data collectors using standardized instructions to ensure consistency in questionnaire administration. Participants were assured of confidentiality and informed that their responses would not affect their clinical care, thereby minimizing social desirability bias. Potential confounding by age, gender, and duration of diabetes was addressed analytically by including these variables in multivariable models, given their established associations with diabetes-related knowledge and self-care behaviors (18).

The sample size was set at 180 participants to estimate KAP levels with acceptable precision, assuming moderate prevalence of adequate knowledge and good practices, a 95% confidence level, and a feasible recruitment target within the study period. Data were entered into a secure, password-protected database and double-checked for accuracy. Statistical analysis was performed using SPSS (IBM Corp., Armonk, NY). Categorical variables were summarized as frequencies and percentages, while continuous variables were summarized as means and standard deviations. Associations between KAP domains and participant characteristics were examined using chi-square tests, and multivariable logistic regression analysis was conducted to identify independent predictors of good self-care practices. A two-sided p-value of <0.05 was considered statistically significant. Ethical approval for the study was obtained from the relevant institutional ethics review committee in Lahore, and all procedures were conducted in accordance with the principles outlined in the Declaration of Helsinki for research involving human participants (19). Data integrity and reproducibility were ensured through predefined variable coding, standardized data collection procedures, and restricted access to anonymized study data limited to the research team.

RESULTS

A total of 180 patients attending primary health care facilities in Lahore were included in the analysis. Slightly more than half of the participants were male (54.4%), and the largest proportion belonged to the 31–50 years age group (46.1%). One-quarter of the participants were older than 50 years, while 28.9% were aged between 18 and 30 years. More than half of the participants (58.3%) reported a duration of diabetes exceeding five years. Regarding educational status, 35.6% had primary education or below, 39.4% had secondary education, and 25.0% had attained higher education (Table 1). Overall, adequate knowledge regarding diabetes mellitus was observed in 54.4% of participants, while 62.8% demonstrated a positive attitude toward diabetes management. In contrast, only 41.7% of participants reported good self-care practices, indicating a substantial gap between knowledge, attitudes, and actual behaviors (Table 2). Analysis of factors associated with adequate knowledge revealed that educational level was a significant determinant. Participants with higher education had a markedly greater likelihood of possessing adequate knowledge compared with those with secondary or lower education (75.6% vs 47.4%; OR 3.12; 95% CI 1.46–6.66; $p=0.003$). No statistically significant associations were observed between adequate knowledge and gender or age group (Table 3). A strong relationship was observed between knowledge, attitude, and self-care practices. Participants with adequate knowledge were more than three times as likely to report good practices compared with those with inadequate knowledge (53.1% vs 28.0%; OR 3.47; 95% CI 1.84–6.55; $p<0.001$). Similarly, a positive attitude toward diabetes management was strongly associated with good practices, with over half of participants with positive attitudes demonstrating good practices compared with 25.4% among those with negative or neutral attitudes (OR 4.18; 95% CI 2.17–8.05; $p<0.001$) (Table 4). Evaluation of specific self-care behaviors showed that medication adherence was the most commonly reported practice, with 68.9% of participants reporting regular adherence. In contrast, adherence to recommended diabetic diets (39.4%) and regular physical activity (36.1%) were notably low. Routine blood glucose monitoring was practiced by 44.4% of participants, while regular follow-up visits were reported by 51.1% (Table 5). In multivariable logistic regression analysis, adequate knowledge (adjusted OR 2.96; 95% CI 1.49–5.89; $p=0.002$), positive attitude (adjusted OR 3.41; 95% CI 1.72–6.77; $p<0.001$), and higher educational attainment (adjusted OR 2.18; 95% CI 1.07–4.45; $p=0.031$) remained independent predictors of good self-care practices after adjustment for age and gender. Duration of diabetes and age were not independently associated with good practices in the adjusted model (Table 6).

Table 1. Socio-Demographic Characteristics of Participants (n = 180)

Variable	Category	n (%)
Gender	Male	98 (54.4)
	Female	82 (45.6)
Age group (years)	18–30	52 (28.9)
	31–50	83 (46.1)

Duration of diabetes	>50	45 (25.0)
	≤5 years	75 (41.7)
	>5 years	105 (58.3)
Education level	Primary or below	64 (35.6)
	Secondary	71 (39.4)
	Higher	45 (25.0)

Table 2. Overall, Knowledge, Attitude, and Practice Levels (KAP) (n = 180)

Domain	Category	n (%)
Knowledge	Adequate	98 (54.4)
	Inadequate	82 (45.6)
Attitude	Positive	113 (62.8)
	Negative/Neutral	67 (37.2)
Practices	Good	75 (41.7)
	Poor	105 (58.3)

Table 3. Association of Socio-Demographic Factors with Adequate Knowledge (n = 180)

Variable	Category	Adequate knowledge n/N (%)	Odds Ratio (95% CI)	p-value
Gender	Male	58/98 (59.2)	1.43 (0.79–2.61)	0.232
	Female	40/82 (48.8)	Reference	—
Age group	≤50 years	74/135 (54.8)	1.07 (0.56–2.05)	0.838
	>50 years	24/45 (53.3)	Reference	—
Education	Higher	34/45 (75.6)	3.12 (1.46–6.66)	0.003
	≤Secondary	64/135 (47.4)	Reference	—

Table 4. Association of Knowledge and Attitude with Good Practices (n = 180)

Variable	Category	Good practices n/N (%)	Odds Ratio (95% CI)	p-value
Knowledge	Adequate	52/98 (53.1)	3.47 (1.84–6.55)	<0.001
	Inadequate	23/82 (28.0)	Reference	—
Attitude	Positive	58/113 (51.3)	4.18 (2.17–8.05)	<0.001
	Negative/Neutral	17/67 (25.4)	Reference	—

Table 5. Diabetes Self-Care Practices by Specific Behaviors (n = 180)

Practice	Yes n (%)	No n (%)
Regular medication adherence	124 (68.9)	56 (31.1)
Adherence to diabetic diet	71 (39.4)	109 (60.6)
Regular physical activity	65 (36.1)	115 (63.9)
Routine blood glucose monitoring	80 (44.4)	100 (55.6)
Regular follow-up visits	92 (51.1)	88 (48.9)

Table 6. Multivariable Logistic Regression Analysis: Predictors of Good Practices (n = 180)

Predictor	Adjusted OR	95% CI	p-value
Adequate knowledge	2.96	1.49–5.89	0.002
Positive attitude	3.41	1.72–6.77	<0.001
Higher education	2.18	1.07–4.45	0.031
Duration of diabetes >5 years	1.29	0.67–2.49	0.441
Age >50 years	1.11	0.54–2.28	0.771

DISCUSSION

This cross-sectional study provides insight into the levels of knowledge, attitude, and practices regarding diabetes mellitus among patients attending primary health care settings in Lahore. The findings indicate that while more than half of the participants demonstrated adequate knowledge and nearly two-thirds exhibited a positive attitude toward diabetes management, fewer than half reported good self-care practices. This discrepancy highlights a critical gap between awareness and actual behavior, a pattern that has been consistently reported in diabetes-related KAP studies across diverse populations (20,21).

The proportion of participants with adequate knowledge observed in this study is comparable to findings from similar studies conducted in urban primary care settings in South Asia, where moderate levels of diabetes awareness have been documented (22). Higher educational attainment emerged as a significant determinant of

adequate knowledge, underscoring the role of formal education in facilitating understanding of chronic disease processes and management principles. This association aligns with existing evidence suggesting that education enhances health literacy, which in turn influences patients' ability to comprehend medical advice and engage in self-care behaviors (23).

Attitudes toward diabetes management were generally positive among participants, with a majority expressing belief in the controllability of diabetes through treatment and lifestyle modification. Positive attitudes were strongly associated with good self-care practices, even after adjustment for potential confounders. This finding supports behavioral models that emphasize the role of individual perceptions, motivation, and self-efficacy in translating knowledge into action (24). Patients who perceive diabetes as manageable and believe in the effectiveness of lifestyle changes are more likely to adhere to recommended practices.

Despite relatively favorable knowledge and attitudes, self-care practices were suboptimal, particularly in relation to dietary adherence and physical activity. While medication adherence was reported by nearly two-thirds of participants, fewer than 40% followed recommended dietary guidelines or engaged in regular physical activity. Similar gaps have been reported in previous studies, suggesting that lifestyle modification remains one of the most challenging aspects of diabetes management (25). Cultural dietary preferences, limited access to structured counseling, time constraints, and lack of supportive environments for physical activity may contribute to poor adherence to lifestyle recommendations in primary care populations (26).

The strong independent association between adequate knowledge, positive attitude, and good practices observed in the multivariable analysis highlights the interrelated nature of KAP domains. However, the persistence of poor practices among a substantial proportion of knowledgeable patients indicates that information alone is insufficient to change behavior. This underscores the need for comprehensive, behaviorally informed interventions that go beyond knowledge dissemination to address motivational, social, and structural barriers to self-care (27).

Primary health care settings represent a critical platform for implementing such interventions, as they provide ongoing contact with patients and opportunities for repeated counseling and follow-up. Integrating structured diabetes education programs, personalized goal-setting, and culturally appropriate lifestyle counseling into routine primary care services may enhance self-care practices and improve long-term outcomes (28). Training primary health care providers in patient-centered communication and behavioral change techniques may further strengthen diabetes management at this level of care.

Several limitations should be considered when interpreting these findings. The cross-sectional design precludes causal inference, and the use of convenience sampling may limit generalizability beyond the study population. Additionally, self-reported practices may be subject to recall and social desirability bias. Nevertheless, the study provides valuable local evidence on diabetes-related knowledge, attitudes, and practices among primary care patients in Lahore and addresses an important gap in the existing literature.

Overall, the findings emphasize that improving diabetes outcomes in primary health care settings requires a multifaceted approach that combines patient education with strategies to foster positive attitudes and facilitate sustainable behavior change, particularly in lifestyle-related practices (29).

CONCLUSION

This study demonstrates that while a substantial proportion of patients attending primary health care settings in Lahore possess adequate knowledge and positive attitudes toward diabetes mellitus, self-care practices remain insufficient. Adequate knowledge, positive attitude, and higher educational attainment were identified as key predictors of good diabetes-related practices, highlighting the interconnected nature of these domains. The observed gap between awareness and behavior underscores the need for comprehensive diabetes management strategies that extend beyond information provision to address behavioral and contextual barriers. Strengthening structured diabetes education, continuous counseling, and patient-centered support within primary health care services may play a critical role in improving self-care practices and reducing the risk of diabetes-related complications.

REFERENCES

1. World Health Organization. Diabetes. Geneva: World Health Organization; 2023.
2. International Diabetes Federation. IDF Diabetes Atlas. 10th ed. Brussels: International Diabetes Federation; 2021.
3. Powers MA, Bardsley J, Cypress M, Duker P, Funnell MM, Fischl AH, et al. Diabetes self-management education and support in adults with type 2 diabetes. *Diabetes Care*. 2020;43(7):1636–49.

4. Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM. Self-management education for adults with type 2 diabetes. *Diabetes Care*. 2002;25(7):1159–71.
5. Shrivastava SR, Shrivastava PS, Ramasamy J. Role of self-care in management of diabetes mellitus. *J Diabetes Metab Disord*. 2013;12:14.
6. Al-Qazaz HK, Sulaiman SA, Hassali MA, Shafie AA, Sundram S, Al-Nuri R, et al. Diabetes knowledge, medication adherence and glycemic control among patients with type 2 diabetes. *Int J Clin Pharm*. 2011;33(6):1028–35.
7. Basit A, Fawwad A, Qureshi H, Shera AS. Prevalence of diabetes, pre-diabetes and associated risk factors: Second National Diabetes Survey of Pakistan. *BMJ Open*. 2018;8(8):e020961.
8. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, et al. Global and regional diabetes prevalence estimates for 2019. *Diabetes Res Clin Pract*. 2019;157:107843.
9. Afroz A, Ali L, Karim MN, Alramadan MJ, Alam K, Magliano DJ, et al. Glycaemic control for people with type 2 diabetes in Bangladesh. *Diabetes Res Clin Pract*. 2019;152:166–77.
10. Murugesan N, Snehathatha C, Shobhana R, Roglic G, Ramachandran A. Awareness about diabetes and its complications in the general and diabetic population. *Diabetes Res Clin Pract*. 2007;77(3):433–7.
11. Saleh F, Mumu SJ, Ara F, Begum HA, Ali L. Knowledge, attitude and practice of type 2 diabetic patients regarding diabetes and its complications. *Mymensingh Med J*. 2012;21(3):418–25.
12. Riaz M, Basit A, Fawwad A, Ahmadani MY. Diabetes awareness and practices in Pakistan. *J Pak Med Assoc*. 2016;66(4):486–9.
13. Islam FMA, Chakrabarti R, Dirani M, Islam MT, Ormsby G, Wahab M, et al. Knowledge, attitudes and practice of diabetes in rural Bangladesh. *Diabetes Res Clin Pract*. 2014;105(3):e7–10.
14. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The STROBE statement. *PLoS Med*. 2007;4(10):e296.
15. Fitzgerald JT, Funnell MM, Hess GE, Barr PA, Anderson RM, Hiss RG, et al. The reliability and validity of a brief diabetes knowledge test. *Diabetes Care*. 1998;21(5):706–10.
16. Anderson RM, Funnell MM, Fitzgerald JT, Marrero DG. The diabetes empowerment scale. *Diabetes Care*. 2000;23(6):739–43.
17. Kassahun T, Gesesew H, Mwanri L, Eshetie T. Diabetes related knowledge, self-care behaviours and adherence. *BMC Endocr Disord*. 2016;16:28.
18. Walker RJ, Smalls BL, Hernandez-Tejada MA, Campbell JA, Egede LE. Effect of diabetes self-efficacy on glycemic control. *Endocrine*. 2014;47(2):571–8.
19. World Medical Association. Declaration of Helsinki. *JAMA*. 2013;310(20):2191–4.
20. Bukhsh A, Khan TM, Nawaz MS, Ahmed HS, Chan KG, Lee LH, et al. Association of diabetes knowledge with glycemic control. *Patient Prefer Adherence*. 2019;13:19–28.
21. Kugbey N, Takyi J, Adulai K. Knowledge, attitude and practice of diabetes. *BMC Public Health*. 2017;17:862.
22. Gulabani M, John M, Isaac R. Knowledge of diabetes, its treatment and complications. *Indian J Community Med*. 2008;33(3):204–6.
23. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes. *Ann Intern Med*. 2011;155(2):97–107.
24. Bandura A. Health promotion by social cognitive means. *Health Educ Behav*. 2004;31(2):143–64.
25. Khattab M, Khader YS, Al-Khawaldeh A, Ajlouni K. Factors associated with poor glycemic control. *J Diabetes Complications*. 2010;24(2):84–9.
26. Hu FB. Globalization of diabetes. *Diabetes Care*. 2011;34(6):1249–57.
27. Glasgow RE, Peeples M, Skovlund SE. Where is the patient in diabetes performance measures? *Diabetes Care*. 2008;31(5):1046–50.
28. Chryala CA, Sherr D, Lipman RD. Diabetes self-management education for adults with type 2 diabetes. *Patient Educ Couns*. 2016;99(6):926–43.
29. Fisher EB, Boothroyd RI, Elstad EA, Hays L, Henes A, Maslow GR, et al. Peer support of complex health behaviors. *Am J Prev Med*. 2012;42(5):S128–34.